

Fairbanks Fishery Resources Progress Report Number FY86-6

THE FRESHWATER FOOD HABITS OF JUVENILE
ARCTIC CHAR IN STREAMS IN THE ARCTIC NATIONAL WILDLIFE REFUGE, ALASKA

Key Words

Food habits, Arctic char, Arctic National Wildlife Refuge

Tonya M. Stevens
Stephen J. Deschermeier

Fairbanks Fishery Resources Station
U.S. Fish and Wildlife Service
101 12th Avenue, Box 20, Rm. 110
Fairbanks, Alaska 99701

24 September 1986

TABLE OF CONTENTS

	<u>Page</u>
List of tables	i
List of figures	ii
Abstract	iii
Introduction	1
Methods	1
Results	8
Discussion	8
Literature Cited	10

LIST OF TABLES

	<u>Page</u>
1. Dates of fish sampling, gear type used, fork length range of fish collected, number of Arctic char stomachs collected and the number of empty stomachs, August 1982 - July 1985, Arctic National Wildlife Refuge, Alaska	3
2. Stomach contents of juvenile Arctic char, by drainage, from four drainages of the Arctic National Wildlife Refuge, Alaska, 1982 - 1985, arranged by drainage.	5
3. Stomach contents of juvenile Arctic char, by length group, from four drainages of the Arctic National Wildlife Refuge, Alaska, 1982 - 1985, arranged by size class of fish.	7

LIST OF FIGURES

	<u>Page</u>
1. Collection sites on four river drainages in the Arctic National Wildlife Refuge, Alaska	2

The freshwater food habits of juvenile Arctic char in streams in the Arctic National Wildlife Refuge, Alaska.

Tonya M. Stevens and Stephen J. Deschermeier. U.S. Fish and Wildlife Service, Fishery Resources, Fairbanks, Alaska.

Abstract:

Juvenile Arctic char (Salvelinus alpinus) were collected from the Canning, Hulahula, Aichilik, and Kongakut Rivers in the Arctic National Wildlife Refuge in Alaska between 1982 and 1985. Stomachs were removed from 189 fish and examined for content. Chironomids were the most commonly found prey organism. Plecopterans were second overall except in samples from the Canning River in which Ephemeropterans were more commonly found.

The freshwater food habits of juvenile Arctic char in streams in the Arctic National Wildlife Refuge, Alaska.

Tonya M. Stevens and Stephen J. Deschermeier. U.S. Fish and Wildlife Service, Fishery Resources, Fairbanks, Alaska.

Abstract:

Juvenile Arctic char (Salvelinus alpinus) were collected from the Canning, Hulahula, Aichilik, and Kongakut Rivers in the Arctic National Wildlife Refuge in Alaska between 1982 and 1985. Stomachs were removed from 189 fish and examined for content. Chironomids were the most commonly found prey organism. Plecopterans were second overall except in samples from the Canning River in which Ephemeropterans were more commonly found.

INTRODUCTION

The Fairbanks Fishery Resources Office has conducted baseline survey and inventory studies of the fishery resources of the North Slope of the Arctic National Wildlife Refuge (ANWR) since 1981 under the direction of Sec. 1002(c) of the Alaska National Interest Lands Conservation Act of 1980. Arctic char (*Salvelinus alpinus*) were sacrificed during these studies for age and growth information and digestive tracts were often removed to learn food habits of fish in the study streams. Some of the char collected could not be aged; however, it is assumed that all char examined for this report were juveniles (under 5 years), based on length comparisons with fish of known ages taken from the same streams at the same time of year.

Four life history types have been described for North Slope Arctic char (McCart 1980): one anadromous form and 3 non-anadromous forms. The non-anadromous forms have been categorized as stream-residents [stream-resident char have also been referred to as "residual Arctic char" (Craig 1977a)], lake-residents or spring-residents. The juveniles examined for this report were either anadromous, stream-resident, or spring-resident char; due to the difficulty of differentiating between these types during the juvenile stage, only those char taken from Shublik Spring (a tributary to the Canning River) are of known type. No separation between stream-resident and anadromous juveniles was made for this analysis.

The objective of this study was to provide baseline information concerning food habits of juvenile Arctic char from streams on the North Slope of the ANWR.

METHODS

Juvenile Arctic char were taken from the Canning River in 1982, from the Aichilik River in 1982 and 1983, from the Hulahula River in 1983, and from the Kongakut River in 1985 (Fig. 1). The sampling sites are described in detail in Smith and Glesne (1983), Daum et al. (1984), and Deschermeier et al. (1986). Fish were collected with a variety of gear: hook and line, electroshocker, minnow traps, gillnets, and seines (Table 1). Table 1 gives the date and location of the sampling sites on each river, the sample sizes, the number of fish collected, and the number of empty stomachs.

Both whole fish and separate digestive tracts were preserved in the field in solutions of approximately 10% formalin. The fork lengths of whole fish measured after preservation were adjusted for shrinkage (Parker 1963). Stomach contents were removed from formalin and transferred to isopropyl alcohol before examination.

Keys used for identification of organisms comprising stomach contents were Oliver et al. (1978), Pennak (1978), and Merritt and Cummins (1984). Organisms were viewed with a 1-7x Bausch and Lomb dissecting scope and a Bausch and Lomb microscope (10-40x).

Sample sizes were insufficient for statistical testing, therefore stomach contents were not analyzed in detail. Relative biomass of food organisms was




Figure 1. Collection sites on four river drainages in the Arctic National Wildlife Refuge, Alaska. Table 1 gives sampling dates and location names for each number on the map.

Table 1. Dates of fish sampling, gear type used, fork length range of fish collected, number of Arctic char stomachs collected and the number of empty stomachs, August 1982 - July 1985, Arctic National Wildlife Refuge, Alaska; ES = electroshocker, MT = baited minnow traps, S = seine, GN = gill-net, A = angling.

Figure 1 Location	Sampling site no.	Sampling date	Gear type	Fork length range (mm)	Number of stomachs collected	Number of empty stomachs
<u>Canning River</u>						
Coastal trib.	1	8-4-82	ES	57 - 155	7	0
Coastal trib.	2	8-2-82	ES	83 - 151	15	0
		8-9-82	MT	77 - 136	4	0
		8-10-82	MT	85 - 106	2	0
		8-11-82	ES	60 - 136	3	0
		8-12-82	MT	135	1	0
		8-14-82	MT	128 - 140	3	1
Shublik Spring	3	9-19-82	ES	45 - 158	9 <u>44</u>	0 <u>1</u>
<u>Hulahula River</u>						
Near mouth	1	8-8-83	S	64	1	0
	1	8-31-83	S	66 - 72	4	1
Lower river	2	7-16-83	ES	78	1	0
Old Man Creek	3	7-16-83	ES	65 - 110	13	0
Fish hole II	4	8-15-83	ES	45 - 93	3	0
	4	9-19-83	MT	51 - 131	12	3
Katuk Creek	5	7-16-83	ES	69 - 136	13	0
W. Patuk Creek	6	7-15-83	ES	106 - 185	22	1
E. Patuk Creek	7	7-15-83	ES	99 - 136	7	1
Grasser Strip	8	8-13-83	ES	174 - 218	2 <u>78</u>	0 <u>6</u>

Table 1 continued

Figure 1 Location	Sampling date	Gear type	Fork length range (mm)	Number stomachs collected	Number stomachs empty
<u>Aichilik River</u>					
1	7-4-83	ES	117 - 178	7	0
2	7-3-83	ES	118 - 182	5	0
3	7-4-83	ES	74 - 171	15	8
4	9-20-82	MT GN	117 - 131 125	2 1	0 0
	9-21-82	A MT GN	107 70 - 108 113 - 120	1 3 2 36	0 0 0 8
<u>Kongakut River</u>					
1	7-2-85	MT	73 - 153	18	5
2	7-1-85	MT	86 - 141	10	3
3	6-29-85	MT	149 - 160	2	0
4	6-24-85	MT	101	1 31	0 8
Totals				189	23

Table 2. Stomach contents of juvenile Arctic char from four drainages of the Arctic National Wildlife Refuge, Alaska, 1982 - 1985; A = adult, P = pupae, L = larvae, N = nymph. Results are given as percent frequency of occurrence and percent composition of stomach items. Empty stomachs were not included in the calculations.

NOT REPRODUCED IN THIS DOCUMENT

Organism		Canning River		Hulahula River		Aichilik River		Kongakut River	
		Freq.	Comp.	Freq.	Comp.	Freq.	Comp.	Freq.	Comp.
Unidentified Diptera	A	48.8	2.6	50.0	0.7	53.6	0.8	12.5	0.6
	P	16.3	1.3	11.0	0.2	17.9	0.1	12.5	0.6
	L			0.5	*	3.6	*	4.2	0.2
Chironomidae	P	58.1	8.5	29.2	1.0	53.6	1.3	8.3	0.4
	L	93.0	74.6	98.6	92.0	92.9	88.0	91.7	86.0
Tipulidae	L	4.7	0.1	27.8	*	10.7	*	41.2	3.6
Simuliidae	P	18.6	0.5						
	L	30.2	2.9	15.3	0.2	28.6	1.4		
Empididae	L	9.3	0.1	22.2	*	3.6	*	16.7	2.1
Psychodidae	L	2.3	0.1	1.4	*				
Plecoptera	N	23.3	0.5	62.5	5.0	53.6	2.3	29.2	2.3
	N	55.8	6.5	27.8	0.6	28.6	3.9	16.7	1.3
Coleoptera	A	13.9	0.3	6.9	*	17.9	0.1	8.3	0.6
	L	7.0	0.1	6.9	*	25.0	0.6	4.2	*
Trichoptera	L	25.6	1.3	2.8	*				
Hemiptera	A			9.7	*	14.3	0.1		
Collembola	A	2.3	*	6.9	*	7.1	*		

Table 2 continued

Organism	Canning River		Hulahula River		Aichilik River		Kongakut River	
	Freq.	Comp. %	Freq.	Comp. %	Freq.	Comp. %	Freq.	Comp. %
Amphipoda					7.1	1.3		
Hydracarina	16.3	0.3	1.4	*	3.6	*	4.2	0.2
Araneae			6.9	*	7.1	*		
Nematoda	9.3	0.1	13.9	0.1	14.3	0.1	4.2	0.2
Unidentified terrestrial insects	9.3	0.1	19.4	0.1	10.7	*		
Fish	2.3	n	2.8	n			16.7	n
Sample size	43		73		26		24	
Total number of organisms		6,207		42,874		23,438		472

* = Items that constituted less than 0.1 % of the composition.
n - Fish were not included in calculation of percent composition.

Table 3. Stomach contents of juvenile Arctic char, by length group, from four drainages in the Arctic National Wildlife Refuge, Alaska, 1982 - 1985. Results are given as percent frequency of occurrence and percent composition of stomach items. Fish captured in minnow traps and those with empty stomachs were not included in the analysis.

Organism		Size class					
		40 - 99 mm		100 - 149 mm		150 - 221 mm	
		% Freq.	% Comp.	% Freq.	% Comp.	% Freq.	% Comp.
Unidentified							
Diptera	A	30.8	0.6	62.1	0.7	70.8	1.0
	P	15.4	0.3	12.1	0.4	4.2	*
	L			1.7	*	16.7	*
Chironomidae	P	26.9	1.5	39.7	0.7	54.2	3.4
	L	88.5	80.2	98.2	93.0	95.8	86.2
Tipulidae	L	7.7	*	29.3	0.1	16.7	*
Simuliidae	P	3.8	0.1	5.2	*	4.2	*
	L	23.1	3.3	24.1	0.5	25.0	0.7
Empididae	L	3.8		27.6	0.1	8.3	
Psychodidae	L					4.2	*
Plecoptera	N	38.5	9.3	53.4	2.0	54.2	4.9
Emphemeroptera	N	61.5	4.2	17.2	1.4	41.7	2.5
Coleoptera	A			13.	*	20.8	0.1
	L	3.8	*	10.3	*	37.5	0.5
Trichoptera	L	3.8	0.1	10.3	0.1	16.7	0.2
Hemiptera	A			6.9	*	20.8	0.1
Collembola	A	3.8	*	5.2	*	16.7	*
Amphipoda				3.4	0.7	4.2	*
Hydracarina				5.2	*	29.2	0.1
Araneae		3.8	*	3.4	*	16.7	*
Nematoda		7.7	0.1	10.3	0.1	25.0	0.2
Unidentified terrestrial insects		3.8	0.1	20.7	0.1	33.3	0.1
Fish				3.4	n	4.2	n
Sample size		26		58		24	
Total organisms			4,296		43,694		27,312

* = Items that constituted less than 0.1 % of the stomach contents.
n - Fish were not included in the calculation of percent composition

not taken into account, nor was there an attempt to identify the contribution of micro-organisms, detritus or plant material in the diet of the juvenile Arctic char examined for this report.

Stomach contents are listed in terms of percent frequency of occurrence and percent composition of all stomachs combined. In Table 2, results are grouped by the drainage from which fish were taken; in Table 3, results are grouped by arbitrarily determined size classes.

RESULTS

Dipteran larvae of the family Chironomidae comprised the overwhelming majority of the diet of almost all fish examined, in terms of both frequency of occurrence and percent composition. Only 2 of 166 fish examined contained other organisms in greater number; these two char (from the lower Aichilik River) were full of amphipods. Adult dipterans also occurred frequently but comprised a small proportion of the diet. The frequency of occurrence of dipteran adults increased with an increase in size class.

The next most frequently eaten organisms appeared to be Plecopteran (stonefly) nymphs, except in the fish from the Canning River, in which Ephemeropteran (mayfly) nymphs were found more frequently and made up a greater portion of the total contents.

Seven char from three drainages had remains of fish in their stomachs. These remains were presumed to be Arctic grayling (Thymallus arcticus) young-of-the-year, based on their sizes. The smallest char, from the Kongakut River, was 118 mm long and weighed 10 g, and had recently eaten 2 grayling each 18 mm long.

The stomach contents of the spring-resident char differed slightly from those of fish from the main Canning River. They contained many Trichoptera larvae and larvae of the dipteran family Psychodidae. These organisms were found only in stomachs of fish from Shublik Spring and in a few fish from the Hulahula River.

DISCUSSION

The stomach contents of juvenile Arctic char from four drainages flowing into the Beaufort Sea appear to be similar to those examined in other studies (Bain 1974, Craig 1977a, b, c). Dipteran larvae, primarily Chironomids, constituted the greatest portion of the diet.

Although these char appeared to rely greatly on Chironomid larvae as a food source, there were many other insect and invertebrate orders represented in stomach samples. It was not difficult to judge that Chironomids dominated in both numbers and biomass; however, the contribution of other organisms to the diet of juvenile char is not apparent when merely reviewing the tables presented in this report. McCart (1980) discussed the inadequacies of describing food habits using data expressed in the form of percent frequency of occurrence and percent composition. Despite the drawbacks to this system,

it was chosen as the style of analysis and presentation because it has been used by other researchers who have reported char food habits (Grainger 1953, Bain 1974, Craig 1977a, b, c, Sparholt 1985) and thereby allows gross comparisons to be made between fish from different areas.

Most unfairly represented in terms of contribution to the diet were the stonefly and mayfly nymphs which were found in the majority of stomachs. These organisms varied greatly in size ; many of the char stomachs contained large nymphs in small numbers. The data describing these stomach contents showed that whereas either mayfly or stonefly nymphs may have occurred quite frequently, they comprised a small percent of the numbers of organisms composing the diet in general. This should not mislead readers to believe that they were an unimportant component of the diet. Most other organisms were more satisfactorily represented in Tables 2 and 3 than stonefly and mayfly nymphs.

Ephemeropterans were found in greater frequency than plecopterans only in stomachs of the fish from the Canning River. The difference is most likely due to the varying time of emergence of the nymphs in the areas in question, rather than an actual preference of char for mayfly nymphs on the Canning.

Johnson (1980) and McCart (1980) describe char as opportunistic feeders. The variety of organisms and the differences in composition of stomach contents of char from different drainages in this study are indications that their classification is likely true for the juvenile char from streams of the North Slope of the Arctic National Wildlife Refuge.

LITERATURE CITED

- Bain, L. H. 1974. Life histories and systematics of arctic char (Salvelinus alpinus, L.) in the Babbage River System, Yukon Territory. Chap. 1, pages 1-156 in P.J. McCart, ed. Life histories of three species of freshwater fishes in Beaufort Sea drainages, Yukon Territory, Can. Arctic Gas Study Ltd. Bio. Rep. Ser. 18.
- Craig, P.C. 1977a. Ecological studies of anadromous and resident populations of Arctic char in the Canning River drainage and adjacent waters of the Beaufort Sea, Alaska. Chap. 1, pages 1 - 116 in P.J. McCart, ed. Fisheries investigations along the North Slope and the Beaufort Sea coast in Alaska with emphasis on Arctic char. Can. Arctic Gas Study Ltd. Bio. Rep. Ser. 41.
- Craig P.C. 1977b. Arctic char in Sadlerochit Spring, Arctic National Wildlife Range. Chap. 2, pages 1-29 in P.J. McCart, ed. Fisheries investigations along the North Slope and the Beaufort Sea coast in Alaska with emphasis on Arctic char. Can. Arctic Gas Study Ltd. Bio. Rep. Ser. 41.
- Craig, P. C. 1977c. Fisheries investigations in the Shaviovic River drainage with emphasis on Arctic char in the Kavik River. Chap. 3, pages 1-28 in P.J. McCart, ed. Fisheries investigations along the North Slope and Beaufort Sea Coast in Alaska with emphasis on Arctic char. Can. Arctic Gas Study Ltd. Bio. Rep. Ser. 41.
- Daum, D., P. Rost, and M. Smith. 1984. Fisheries studies on the North Slope of the Arctic National Wildlife Refuge, 1983. Pages 464-522 in G.W. Garner and P.E. Reynolds, eds. 1983 update report baseline study of the fish, wildlife and their habitats. U.S. Fish and Wildl. Serv., Anchorage, Ak. 614 pp.
- Deschermeier, S. J., T. Stevens, D. Wiswar, and R. West. 1986. Fisheries investigations on the Kongakut River, Arctic National Wildlife Refuge, Alaska 1985. Pages ____ in G.W. Garner and P.E. Reynolds, eds. 1986 update report baseline study of the fish, wildlife and their habitats. U.S. Fish and Wildl. Serv., Anchorage, Ak.
- Glesne, R. and S. J. Deschermeier. 1984. Abundance, distribution and diversity of aquatic macroinvertebrates on the North Slope of the Arctic National Wildlife Refuge, 1982 and 1983. Pages 523-552 in G.W. Garner and P.E. Reynolds, ed. 1983 update report baseline study of the fish, wildlife, and their habitats. U.S. Fish and Wildl. Serv., Anchorage, Ak. 614 pp.
- Grainger, E. H. 1953. On the age, growth, migration, reproductive potential and feeding habits of the arctic char (Salvelinus alpinus) of Frobisher Bay, Baffin Island, Canada. J. Fish. Res. Bd. Can. 10:326-370.
- McCart, P.J. 1980. A review of the systematics and ecology of the Arctic char, Salvelinus alpinus, in the western Arctic. Can. Tech. Rep. Fish. Aquat. Sci. 935.

- Merritt, R.W., and K.W. Cummins, eds. 1984. An introduction to the aquatic insects of North America, second edition. Kendall/Hunt Pub. Co., Dubuque, Iowa. 772 pp.
- Oliver, D.R., D. McClymont, and M.E. Roussel. 1978. A key to some larvae of Chironomidae (Diptera) from the Mackenzie and Porcupine River watersheds. Can. Fish. MAR. Serv. Tech Rep. No. 791.
- Pennak, R.W. 1978. Freshwater invertebrates of the United States, second edition. John Wiley and Sons, New York, NY. 769 pp.
- Parker, R.R. 1963. Effects of formalin on length and weight of fishes. J. Fish Res. Bd. Can. FRBC 20: 144-155.
- Smith, M.W., and R. S. Glesne. 1983. Aquatic studies on the North Slope of the Arctic National Wildlife Refuge 1981 and 1982. Pages 291-364 in G.W. Garner and P.E. Reynolds, eds. 1982 update report baseline study of the fish, wildlife, and their habitats. U.S. Fish and Wildl. Serv., Anchorage, Ak. 379 pp.
- Sparholt, H. 1985. The population, survival, growth, reproduction and food of Arctic charr, Salvelinus alpinus (L.), in four unexploited lakes in Greenland. J. Fish Bio. 26: 313-330.